

Pioglitazone suppresses excessive follicular development in murine preantral follicles.

Nagao S, Baba T, Fujibe Y, Adachi S, **Ikeda K**, Morishita M, Kuno Y, Honnma H, Endo T, Kiya T, Saito T.

J Ovarian Res. 2019 Aug 31;12(1):82.

Androgen potentiates the expression of FSH receptor and supports preantral follicle development in mice.

Fujibe Y, Baba T, Nagao S, Adachi S, **Ikeda K**, Morishita M, Kuno Y, Suzuki M, Mizuuchi M, Honnma H, Endo T, Saito T.

J Ovarian Res. 2019 Apr 4;12(1):31.

Pioglitazone is effective for multiple phenotypes of the Zucker fa/fa rat with polycystic ovary morphology and insulin resistance.

Morishita M, Endo T, Baba T, Kuno Y, **Ikeda K**, Kiya T, Honnma H, Saito T.

J Ovarian Res. 2018 Mar 27;11(1):24.

Long-term treatment with dehydroepiandrosterone may lead to follicular atresia through interaction with anti-Mullerian hormone.

Ikeda K, Baba T, Morishita M, Honnma H, Endo T, Kiya T, Saito T.

J Ovarian Res. 2014 Apr 30;7:46.

Excessive androgen exposure in female-to-male transsexual persons of reproductive age induces hyperplasia of the ovarian cortex and stroma but not polycystic ovary morphology.

Ikeda K, Baba T, Noguchi H, Nagasawa K, Endo T, Kiya T, Saito T.

Hum Reprod. 2013 Feb;28(2):453-61.

Assisted reproductive technique increases the risk of placental polyp.

Baba T, Endo T, **Ikeda K**, Shimizu A, Morishita M, Kuno Y, Honnma H, Kiya T, Ishioka S, Saito T.

Gynecol Endocrinol. 2013 Jun;29(6):611-4.

Weight reduction and pioglitazone ameliorate polycystic ovary syndrome after removal of a Sertoli-stromal cell tumor.

Baba T, Endo T, **Ikeda K**, Shimizu A, Morishita M, Kuno Y, Honnma H, Kiya T, Ishioka S, Saito T.

Int J Womens Health. 2012;4:607-11.

Simultaneous presentation of tubal and primary abdominal pregnancies following clomiphene citrate treatment.

Baba T, Endo T, **Ikeda K**, Takenami N, Shimizu A, Morishita M, Honnma H, Ikeda H, Saito T.

Arch Gynecol Obstet. 2012 Aug;286(2):395-8.

Continuous follicle-stimulating hormone exposure from pituitary adenoma causes periodic follicle recruitment and atresia, which mimics ovarian hyperstimulation syndrome.

Kanaya M, Baba T, Kitajima Y, **Ikeda K**, Shimizu A, Morishita M, Honnma H, Endo T, Saito T.

Int J Womens Health. 2012;4:427-31.